


Optics for Engineers

Chapter 3


Charles A. DiMarzio
Northeastern University

July 2012


Compound Lens and Ray Definitions




Correct Ray




Vertex Plane




Matrix Optics Ray



Ray Definition



Translation




Refraction

Ray Definitions

- Ray Information
 - Straight Line
 - Two Dimensions
 - Slope and Intercept
- Mathematical Formulation
 - Column Vector
 - Two Elements
 - Intercept on Top
 - Reference to Local z
 - Angle on Bottom

$$\mathbf{V} = \begin{pmatrix} x \\ \alpha \end{pmatrix}$$

- Some Books Differ



- Arbitrary Operation

$$\mathcal{M} = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$$

$$\mathbf{V}_{end} = \mathcal{M}_{start:end} \mathbf{V}_{start}$$

- Subscript for Vertex Number

Translation From One Surface to the Next

- Move Away from Source
- z_1 to z_2

$$\mathbf{V}_2 = \mathcal{T}_{12} \mathbf{V}_1$$

- Angle Stays Constant


$$\alpha_2 = \alpha_1$$

- Height Changes

$$x_2 = x_1 + \alpha_1 z_{12}$$

- Matrix Form

$$\begin{pmatrix} x_2 \\ \alpha_2 \end{pmatrix} = \begin{pmatrix} 1 & z_{12} \\ 0 & 1 \end{pmatrix} \begin{pmatrix} x_1 \\ \alpha_1 \end{pmatrix}$$



$$\mathcal{T}_{12} = \begin{pmatrix} 1 & z_{12} \\ 0 & 1 \end{pmatrix}$$

Refraction at a Surface (1)

- Matrix Form



$$\mathbf{V}'_1 = \mathcal{R}_1 \mathbf{V}_1$$

- Height Does Not Change

$$x'_1 = (1 \times x_1) + (0 \times \alpha_1)$$

$$\begin{pmatrix} x'_1 \\ \alpha'_1 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ ? & ? \end{pmatrix} \begin{pmatrix} x_1 \\ \alpha_1 \end{pmatrix}$$

- Angle Changes (Ch. 2)



$$\theta = \gamma + \alpha \quad \theta' = \gamma - \beta = \gamma + \alpha'$$

$$\tan \alpha = \frac{p}{s + \delta} \quad \tan \beta = \frac{p}{s' - \delta}$$

$$\tan \gamma = \frac{p}{r - \delta}$$

Refraction at a Surface (2)

- Height Does Not Change

$$\begin{pmatrix} x'_1 \\ \alpha'_1 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ ? & ? \end{pmatrix} \begin{pmatrix} x_1 \\ \alpha_1 \end{pmatrix}$$


- Angle (See Prev. Page)

$$n\theta = n'\theta'$$

$$n(\gamma + \alpha) = n'(\gamma + \alpha'),$$

$$n\frac{x}{r} + n\alpha = n'\frac{x}{r} + n'\alpha'.$$


$$\alpha' = \frac{n - n'}{n'r}x + \frac{n}{n'}\alpha.$$



$$\mathcal{R} = \begin{pmatrix} 1 & 0 \\ \frac{n-n'}{n'r} & \frac{n}{n'} \end{pmatrix}$$

$$\mathcal{R} = \begin{pmatrix} 1 & 0 \\ -\frac{P}{n'} & \frac{n}{n'} \end{pmatrix}$$

Cascading Matrices



$$V_1 = \mathcal{T}_{01} V_0 \quad V_1' = \mathcal{R}_1 V_1 \quad V_2 = \mathcal{T}_{12} V_1' \quad \text{etc.}$$

$$V_{\text{end}} = \mathcal{M}_{0:\text{end}} V_0 \quad \mathcal{M}_{0:\text{end}} = \mathcal{T}_{\text{end}-1:\text{end}} \cdots \mathcal{T}_{12} \mathcal{R}_1 \mathcal{T}_{01}$$

Multiply from Right to Left as Light Moves from Left to Right.

The Simple Lens (1)

- First Surface

$$\mathbf{V}'_1 = \mathcal{R}_1 \mathbf{V}_1$$

- Translation

$$\mathbf{V}_2 = \mathcal{T}_{12} \mathbf{V}'_1$$


- Second Surface

$$\mathbf{V}'_2 = \mathcal{R}_2 \mathbf{V}_2$$

- Result

$$\mathbf{V}'_2 = \mathcal{L} \mathbf{V}_1$$

$$\mathcal{L} = \mathcal{R}_2 \mathcal{T}_{12} \mathcal{R}_1$$



The Simple Lens (2)

- From Previous Page $\mathcal{L} = \mathcal{R}_2 \mathcal{T}_{12} \mathcal{R}_1$

$$\mathcal{L} = \begin{pmatrix} 1 & 0 \\ -\frac{P_2}{n'_2} & \frac{n'_1}{n'_2} \end{pmatrix} \begin{pmatrix} 1 & z_{12} \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ -\frac{P_1}{n'_1} & \frac{n_1}{n'_1} \end{pmatrix} \quad (n_2 = n'_1)$$

- Strange but Useful Grouping

$$\mathcal{L} = \begin{pmatrix} 1 & 0 \\ -\frac{P_t}{n'_2} & \frac{n_1}{n'_2} \end{pmatrix} + \frac{z_{12}}{n'_1} \begin{pmatrix} -P_1 & n_1 \\ \frac{P_1 P_2}{n'_2} & -P_2 \frac{n_1}{n'_2} \end{pmatrix} \quad (P_t = P_1 + P_2)$$

- Initial: $n_1 = n$, Final: $n'_2 = n'$, Lens: $n'_1 = n_\ell$

$$\mathcal{L} = \begin{pmatrix} 1 & 0 \\ -\frac{P_t}{n'} & \frac{n}{n'} \end{pmatrix} + \frac{z_{12}}{n_\ell} \begin{pmatrix} -P_1 & n \\ \frac{P_1 P_2}{n'} & -P_2 \frac{n}{n'} \end{pmatrix}$$

- n_ℓ implicit in P_1 and P_2 , and thus P_t

The Thin Lens (1)

- The Simple Lens (Previous Page)

$$\mathcal{L} = \begin{pmatrix} 1 & 0 \\ -\frac{P_t}{n'} & \frac{n}{n'} \end{pmatrix} + \frac{z_{12}}{n_\ell} \begin{pmatrix} -P_1 & n \\ \frac{P_1 P_2}{n'} & -P_2 \frac{n}{n'} \end{pmatrix}$$

- Geometric Thickness, z_{12}/n_ℓ , Multiplies Second Term
- Set $z_{12} \rightarrow 0$

$$\mathcal{L} = \begin{pmatrix} 1 & 0 \\ -\frac{P}{n'} & \frac{n}{n'} \end{pmatrix} \quad (\text{Thin Lens})$$

$$P = P_t = P_1 + P_2 \quad \text{Correction Term Vanishes}$$

The Thin Lens (2)

- Thin Lens in terms of Focal Lengths

$$\mathcal{L} = \begin{pmatrix} 1 & 0 \\ -\frac{P}{n'} & \frac{n}{n'} \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ -\frac{1}{f'} & \frac{n}{n'} \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ -\frac{n}{n'f} & \frac{n}{n'} \end{pmatrix}$$

- Front Focal Length: $f = FFL$, Back: $f' = BFL$
- Special but Common Case: Thin Lens in Air

$$\mathcal{L} = \begin{pmatrix} 1 & 0 \\ -P & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ -\frac{1}{f} & 1 \end{pmatrix} \quad (\text{Thin Lens in Air})$$

$$f = f' = \frac{1}{P} \quad FFL = BFL \quad \text{Always True if } n' = n$$

General Problems and the ABCD Matrix

- General Equation

$$\mathbf{V}_{end} = \mathcal{M}_{start:end} \mathbf{V}_{start} \quad \begin{pmatrix} x_{end} \\ \alpha_{end} \end{pmatrix} = \begin{pmatrix} m_{11} & m_{12} \\ m_{21} & m_{22} \end{pmatrix} \begin{pmatrix} x_{start} \\ \alpha_{start} \end{pmatrix}$$

- Determinant Condition (Not Completely Obvious)

$$\det \mathcal{M} = \frac{n}{n'} \quad (\det \mathcal{M} = m_{11}m_{22} - m_{12}m_{21})$$

- Abbe Sine Invariant (or Helmholtz or Lagrange Invariant)


$$n'x'd\alpha' = nx d\alpha$$

Abbe Sine Invariant

- Equation

$$n'x'd\alpha' = nxd\alpha$$

- Alternative Derivations
 - Geometric Optics
 - Energy Conservation (C. 12)
- Lens Example
 - Height Decreases by s'/s
 - Angle Increases by s/s'



- Example: IR Detector
 - Diameter
 $D' = 100\mu\text{m}$
 - Collection Cone
 $FOV'_{1/2} = 30^\circ$
- Telescope Front Lens
 - Diameter
 $D = 20\text{cm}$
 - Max. Field of View

$$FOV_{1/2} = \frac{100 \times 10^{-6}\text{m} \times 30^\circ}{20 \times 10^{-2}\text{m}} = 0.0150^\circ$$

Principal Planes Concept (1)

- Thin Lens

$$\mathcal{L} = \begin{pmatrix} 1 & 0 \\ -\frac{P}{n'} & \frac{n}{n'} \end{pmatrix}$$

- Simple Equation
- Easy Visualization
("High-School Optics")
- Good "First Try"

- Arbitrary Lens
- Vertex to Vertex


$$\mathcal{M}_{VV'} = \begin{pmatrix} m_{11} & m_{12} \\ m_{21} & m_{22} \end{pmatrix}$$

- Possible Simplification


$$\mathcal{M}_{HH'} = \mathcal{T}_{V'H'} \mathcal{M}_{VV} \mathcal{T}_{HV}$$

$$\mathcal{M}_{HH'} = \mathcal{L}$$

- Will It Work?




Principal Planes Concept (2)



Convert a Hard Problem to a Simple one

$$\mathcal{M}_{HH'} = \mathcal{T}_{V'H'} \mathcal{M}_{VV} \mathcal{T}_{HV} \quad \mathcal{M}_{HH'} = \mathcal{L}$$



Useful if a Solution Can Be Found
Very Useful if h and h' Are Not Too Large

Finding the Principal Planes

$$\mathcal{L} = \mathcal{M}_{HH'} = \mathcal{T}_{V'H'} \mathcal{M}_{VV} \mathcal{T}_{HV}$$

$$\begin{pmatrix} 1 & 0 \\ -\frac{P}{n'} & \frac{n}{n'} \end{pmatrix} = \begin{pmatrix} 1 & h' \\ 0 & 1 \end{pmatrix} \begin{pmatrix} m_{11} & m_{12} \\ m_{21} & m_{22} \end{pmatrix} \begin{pmatrix} 1 & h \\ 0 & 1 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 0 \\ -\frac{P}{n'} & \frac{n}{n'} \end{pmatrix} = \begin{pmatrix} m_{11} + m_{21}h' & m_{11}h + m_{12} + m_{21}hh' + m_{22}h' \\ m_{21} & m_{21}h + m_{22} \end{pmatrix}$$

$m_{11} + m_{21}h' = 1$	$m_{11}h + m_{12} + m_{21}hh' + m_{22}h' = 0$
$m_{21} = -\frac{P}{n'}$	$m_{21}h + m_{22} = \frac{n}{n'}$

Three Unknowns: Solution if Determinant Condition Satisfied

$h = \frac{\frac{n}{n'} - m_{22}}{m_{21}}$	Determinant Condition? Yes!
$P = -m_{21}n'$	$h' = \frac{1 - m_{11}}{m_{21}}$

No Assumptions Were Made About \mathcal{M} : This Always Works.


Principal Planes

- Principal Planes are Conjugates of Each Other ($m_{12} = 0$)

$$\begin{pmatrix} x_{H'} \\ \alpha_{H'} \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ -\frac{P}{n'} & \frac{n}{n'} \end{pmatrix} \begin{pmatrix} x_H \\ \alpha_H \end{pmatrix}$$

- Unit Magnification Between Them

$$x_{H'} = x_H$$



Note: Principal Planes May Not Be Accessible


Imaging (We Know The Answer)

- Matrix from Object to Image

$$\mathcal{M}_{SS'} = \mathcal{M}_{H'S'} \mathcal{M}_{HH'} \mathcal{M}_{SH} = \mathcal{T}_{s'} \mathcal{M}_{HH'} \mathcal{T}_s$$

- Conjugate Planes

$$x' = (? \times x) + (0 \times \alpha) \quad \mathcal{M}_{SS'} = \begin{pmatrix} m_{11} & 0 \\ m_{21} & m_{22} \end{pmatrix}$$



Imaging Equation for Compound Lens

$$\mathcal{M}_{SS'} = \begin{pmatrix} 1 & s' \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ -\frac{P}{n'} & \frac{n}{n'} \end{pmatrix} \begin{pmatrix} 1 & s \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 - \frac{s'P}{n'} & s - \frac{ss'P}{n'} + \frac{s'n}{n'} \\ -\frac{P}{n'} & -\frac{sP}{n'} + \frac{n}{n'} \end{pmatrix}$$

- Conjugate Plane Rule: $m_{12} = 0$

$$s - \frac{ss'P}{n'} + \frac{s'n}{n'} = 0$$

$$\frac{n}{s} + \frac{n'}{s'} = P$$

- Measure s and s' from h and h' respectively.

Compound Lens Matrix Results

Magnifications ($mm_\alpha = n'/n$)

$$m = 1 - \frac{s'P}{n'} = 1 - \frac{s'}{n'} \left(\frac{n}{s} + \frac{n'}{s'} \right) = -\frac{ns'}{n's}$$

$$m_\alpha = -\frac{s}{n'} \left(\frac{n}{s} + \frac{n'}{s'} \right) + \frac{n}{n'} = -\frac{s}{s'}$$

Imaging Matrix

$$\mathcal{M}_{SS'} = \begin{pmatrix} m & 0 \\ -\frac{P}{n'} & \frac{n'1}{n m} \end{pmatrix}$$

Thick Lens

- Thick–Lens Equation

$$\mathcal{L} = \begin{pmatrix} 1 & 0 \\ -\frac{P_t}{n'} & \frac{n}{n'} \end{pmatrix} + \frac{z_{12}}{n_\ell} \begin{pmatrix} -P_1 & n \\ \frac{P_1 P_2}{n'} & -P_2 \frac{n}{n'} \end{pmatrix}$$

- Power: $P = -m_{21}n'$

$$P = P_1 + P_2 - \frac{z_{12}}{n_\ell} P_1 P_2 \quad f = \frac{n}{P} \quad f' = \frac{n'}{P}$$

- Principal Planes

$$h = -\frac{n' P_2}{n_\ell P} z_{12} \quad h' = -\frac{n P_1}{n_\ell P} z_{12}$$

Thick Lens in Air: The Thirds Rule for Principal Planes

- Principal Planes and Focal Length

$$f = f' = \frac{1}{P} \quad h = -\frac{1}{n_\ell} \frac{P_2}{P} z_{12} \quad h' = -\frac{1}{n_\ell} \frac{P_1}{P} z_{12}$$

- Principal-Plane Spacing

$$z_{HH'} = z_{12} + h + h' = z_{12} \left(1 - \frac{P_2 + P_1}{n_\ell P} \right)$$

$$P \approx P_1 + P_2 \quad z_{HH'} = z_{12} + h + h' \approx z_{12} \left(1 - \frac{1}{n_\ell} \right)$$

Glass	$n_\ell \approx 1.5$	$z_{HH'} = \frac{z_{12}}{3}$
-------	----------------------	------------------------------

Special Cases

$$h = -\frac{n' P_2}{n_\ell P} z_{12} \quad h' = -\frac{n P_1}{n_\ell P} z_{12}$$

h, h' Negative if P, P_1, P_2 Have Same Signs (Often True)

$h = 0$ if $P_2 = 0$ Convex–Plano or Concave–Plano


$h' = 0$ if $P_1 = 0$ Plano–Convex or Plano–Concave

$h' = h$ if $P_2 = P_1$ Biconvex or Biconcave in Air
and $n' = n$

Example: Biconvex Lens in Air

$$P_1 + P_2 = 10 \text{ diopters, or } f = 10 \text{ cm}$$

Solid=Vertices, Dashed=Principal Planes, Dash-Dot=Focal Planes




“Bending” the Lens


$P_1 + P_2 = 10\text{diopters}$, or $f = 10\text{cm}$

Solid=Vertices, Dashed=Principal Planes, Dash-Dot=Focal Planes

Note “Meniscus” Lenses in Germanium



A. Glass



B. Germanium

Example: Compound Lens Matrix (Two Thin Lenses)



$$\mathcal{M}_{V_1, V'_2} = \mathcal{L}_{V_2, V'_2} \mathcal{T}_{V'_1, V_2} \mathcal{L}_{V_1, V'_1} \quad (\text{Thin Lenses})$$

$$\mathcal{M}_{V_1, V'_2} = \begin{pmatrix} 1 & 0 \\ -\frac{1}{f_2} & 1 \end{pmatrix} \begin{pmatrix} 1 & z_{12} \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ -\frac{1}{f_1} & 1 \end{pmatrix}$$

$$\mathcal{M}_{V_1, V'_2} = \begin{pmatrix} 1 - \frac{z_{12}}{f_1} & z_{12} \\ -\frac{1}{f_2} + \frac{z_{12}}{f_1 f_2} - \frac{1}{f_1} & 1 - \frac{z_{12}}{f_2} \end{pmatrix}$$

Compound Lens Results

- Focal Length (Powers add for small separation)

$$\frac{1}{f} = \frac{1}{f_2} + \frac{1}{f_1} - \frac{z_{12}}{f_1 f_2}$$

- Principal Planes

$$h = \frac{\frac{z_{12}}{f_2}}{-\frac{1}{f_2} + \frac{z_{12}}{f_1 f_2} - \frac{1}{f_1}} = \frac{z_{12} f_1}{z_{12} - f_1 - f_2}$$

$$h' = \frac{\frac{z_{12}}{f_1}}{-\frac{1}{f_2} + \frac{z_{12}}{f_1 f_2} - \frac{1}{f_1}} = \frac{z_{12} f_2}{z_{12} - f_1 - f_2}$$

$$h \rightarrow 0 \quad \text{and} \quad h' \rightarrow 0 \quad \text{if} \quad z_{12} \rightarrow 0$$

Special Case: Afocal

$$z_{12} = f_1 + f_2$$


$$\frac{1}{f} = \frac{f_1}{f_1 f_2} + \frac{f_2}{f_1 f_2} - \frac{z_{12}}{f_1 f_2} = 0.$$

$$m_{21} = 0, \quad \frac{1}{f} = 0, \quad \text{or} \quad f \rightarrow \infty \quad (\text{Afocal})$$

$$h \rightarrow \infty \quad h' \rightarrow \infty$$


Principal Planes are not Very Useful Here.

Example: 2X Magnifier (1)



- We Know How to Do This
 - Object at Front Focus of First Lens
 - Intermediate Image at Infinity
 - Final Image at Back Focus of Second Lens
- But Let's Use Matrix Optics for the Exercise


Example: 2X Magnifier (2)



Lens Vendor Data: Glass=BK7 ($n = 1.515$ at $\lambda = 633\text{nm}$)

Parameter	Label	Value
First Lens Focal Length	f_1	100 mm
First Lens Front Radius (LA1509 Reversed)	r_1	Infinite
First Lens Thickness	$z_{v1,v1'}$	3.6 mm
First Lens Back Radius	r_1'	51.5 mm
First Lens "Back" Focal Length	$f_1 + h_1$	97.6 mm
Lens Spacing	$z_{v1',v2}$	20 mm
Second Lens Focal Length	f_2	200 mm
Second Lens Front Radius (LA1708)	r_2	103.0 mm
Second Lens Thickness	$z_{v2,v2'}$	2.8 mm
Second Lens Back Radius	r_2'	Infinite
Second Lens Back Focal Length	$f_2' + h_2'$	198.2 mm

Example: 2X Magnifier (Thin-Lens Approximation)



$$\frac{1}{f} = \frac{1}{100\text{mm}} + \frac{1}{200\text{mm}} - \frac{20\text{mm}}{100\text{mm} \times 200\text{mm}} \quad f = 71.43\text{mm}$$

$$h = \frac{20\text{mm} \times 100\text{mm}}{20\text{mm} - 100\text{mm} - 200\text{mm}} = -7.14\text{mm}$$

$$h' = \frac{20\text{mm} \times 200\text{mm}}{20\text{mm} - 100\text{mm} - 200\text{mm}} = -14.28\text{mm}$$

$$m = -\frac{s'}{s} = -2 \quad s' = 2s, \quad \text{and} \quad \frac{1}{f} = \frac{1}{s} + \frac{1}{s'} = \frac{1}{s} + \frac{1}{2s}$$

$$s = 3f/2 = 107.1\text{mm} \quad s' = 3f = 214.3\text{mm}$$

Lens Thickness Effects

- Start with Equations for Thin Lenses
- Use Principal Planes in Place of Vertices

$$\mathcal{M}_{H_1, H'_2} = \mathcal{L}_{H_2, H'_2} \mathcal{T}_{H'_1, H_2} \mathcal{L}_{H_1, H'_1}$$

- Same Equation as Thin Lens but Different Meaning

$$\mathcal{M}_{H_1, H'_2} = \begin{pmatrix} 1 & 0 \\ -\frac{1}{f_2} & 1 \end{pmatrix} \begin{pmatrix} 1 & z_{12} \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ -\frac{1}{f_1} & 1 \end{pmatrix}$$

- f_1 from H_1 and f'_1 from H'_1
- f_2 from H_2 and f'_2 from H'_2
- z_{12} from H'_1 to H_2

2X Magnifier, Revisited (1)

- Principal Planes

$$h = \frac{20\text{mm} \times 100\text{mm}}{20\text{mm} - 100\text{mm} - 200\text{mm}} = -7.14\text{mm} \quad H_1 \text{ to } H$$

$$h' = \frac{20\text{mm} \times 200\text{mm}}{20\text{mm} - 100\text{mm} - 200\text{mm}} = -14.28\text{mm} \quad H'_2 \text{ to } H'$$


- Spacing (See Next Page)

$$0.713\text{mm}$$

- Object and Image Distances

$$s = \frac{3f}{2} = 107.1\text{mm} \quad s' = 3f = 214.3\text{mm}$$

2X Magnifier, Revisited (2)



A Suggestion: Global Coordinates

- Notation: $zH1$
 - First Letter: z
 - The Remaining Characters: Plane Name (eg. $H1$)
- Need to Set One Plane as $z = 0$
- Example from the Magnifier
 - $z = 0$ at First Vertex
 - $zH1 = -h_1$
 - $zH = zH1 - h = -h_1 - h$
 - (Text Error: Not $z_H = zH1 - h = h_1 - h$)
 - *etc.*

Telescopes (1)

- Afocal Condition

$$\frac{1}{f} = \frac{1}{f_2} + \frac{1}{f_1} - \frac{z_{12}}{f_1 f_2} = 0 \quad \text{if} \quad z_{12} = f_1 + f_2$$

- Vertex Matrix

$$\mathcal{M}_{V_1, V'_2} = \begin{pmatrix} 1 - \frac{f_1 + f_2}{f_1} & f_1 + f_2 \\ -\frac{1}{f_2} + \frac{f_1 + f_2}{f_1 f_2} - \frac{1}{f_1} & 1 - \frac{f_1 + f_2}{f_2} \end{pmatrix} = \begin{pmatrix} -\frac{f_2}{f_1} & f_1 + f_2 \\ 0 & -\frac{f_1}{f_2} \end{pmatrix}$$

- Imaging Matrix

$$\mathcal{M}_{SS'} = \begin{pmatrix} 1 & s' \\ 0 & 1 \end{pmatrix} \begin{pmatrix} -\frac{f_2}{f_1} & f_1 + f_2 \\ 0 & -\frac{f_1}{f_2} \end{pmatrix} \begin{pmatrix} 1 & s \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} ? & 0 \\ ? & ? \end{pmatrix}$$

Telescopes (2)

$$\mathcal{M}_{SS'} = \begin{pmatrix} -\frac{f_2}{f_1} & -s\frac{f_2}{f_1} + f_1 + f_2 - s'\frac{f_1}{f_2} \\ 0 & -\frac{f_1}{f_2} \end{pmatrix} = \begin{pmatrix} ? & 0 \\ ? & ? \end{pmatrix}$$

$$-s\frac{f_2}{f_1} + f_1 + f_2 - s'\frac{f_1}{f_2} = 0$$

$$m = -f_2/f_1, \quad (\text{Afocal})$$


$$ms + f_1 + f_2 + s'/m = 0$$

$$s' = -m^2s - f_1m(1 + m)$$

$$\mathcal{M}_{SS'} = \begin{pmatrix} m & 0 \\ 0 & \frac{1}{m} \end{pmatrix}$$

$$s' \approx -m^2s \quad s \rightarrow \infty$$

Astronomical Telescope



- Magnification: Image is smaller ($\ll 1$) $m = \frac{f_2}{f_1}$
- But a Lot Closer: ($m_z = -m^2$)
- Angular Magnification is Large ($m_\alpha = 1/m$)